

---

## **European Stroke Organization Guidelines for the Management of Intracranial Aneurysms and Subarachnoid Haemorrhage**

Thorsten Steiner<sup>a</sup> Seppo Juvela<sup>d</sup> Andreas Unterberg<sup>b</sup> Carla Jung<sup>b</sup>  
Michael Forsting<sup>c</sup> Gabriel Rinkel<sup>e</sup>

Departments of <sup>a</sup>Neurology and <sup>b</sup>Neurosurgery, Heidelberg University, Heidelberg, and <sup>c</sup>Department of Radiology, University of Essen, Essen, Germany; <sup>d</sup>Department of Clinical Neurosciences, University of Helsinki, Helsinki, Finland; <sup>e</sup>Department Neurology, Utrecht University, Utrecht, The Netherlands

### **Treatment of hyperthermia**

Page 99: Despite treatment with paracetamol to relieve pain, fever develops in more than half of all SAH patients [71], predominantly in those with poor clinical condition on admission and in those with intraventricular extension of the haemorrhage. [71]

Furthermore, fever is an independent risk factor for poor outcome. [69] In around 20% of patients no infection is found, and fever is attributed to inflammatory responses to extravasated blood in the subarachnoid space.

#### **Recommendations**

**Good clinical practice:** Increased temperature should be treated medically and physically.

### **Therapeutic hypothermia**

A randomized trial of mild hypothermia (33 ° C) during aneurysm surgery after SAH showed no benefit. [72]

For full version please click here:

[http://emcools.com/fileadmin/daten/Publikationen/ESO\\_IC\\_Aneurysms\\_SAH\\_Guidelines\\_2013\\_Management\\_of\\_Intracranial\\_Aneurysms\\_and\\_Subarachnoid\\_Haemorrhage.pdf](http://emcools.com/fileadmin/daten/Publikationen/ESO_IC_Aneurysms_SAH_Guidelines_2013_Management_of_Intracranial_Aneurysms_and_Subarachnoid_Haemorrhage.pdf)